



BRUCE M. MacNICOL, DD.S.,M.S.,F.R.CD.(C).
MICHAEL T. JACKSON, B.A., DD.S., M.SC., F.R.CD.(C).
ANDREW P.GATER, H.B.SC., DD.S., F.R.CD.(C).
HURONIA ORAL SURGERY GROUP
Oral and Maxillofacial Surgeons

128 Wellington St. West, Suite 308, Barrie , ON IAN 8J6 Tel (705)
734-9292 Fax (705) 734-0154 www.maxillocare.com

Tooth Exposure Protocol

Your doctor has performed an exposure of an impacted tooth for you. Please note the following.

1. The area is designed to be kept open so don't be concerned if there is some post-operative bleeding from the area. If it is excessive, firm pressure with gauze placed directly on the site for 15 consecutive minutes will get it to resolve.
 - Note: Please be gentle when removing the gauze, particularly if there has been a bracket bonded to the tooth, to avoid pulling the bracket off.
2. If a temporary covering has been placed over the site (called *Barricaid*), don't be concerned when this comes out spontaneously. It is not intended to be over the wound permanently.

If you have any further questions or concerns, please do not hesitate to contact us. We are available at (705) 734-9292.

Thank you.

Huronian Oral Surgery Group